

Lack of genetic structure in greylag goose (*Anser anser*) populations along the European Atlantic flyway

Irene Pellegrino¹, Marco Cucco¹, Arne Follestad² and Mathieu Boos³

¹ University of Piemonte Orientale, DISIT, Alessandria, Italy

² Norsk Institutt for Naturforskning, Trondheim, Norway

³ Naturaconst@, Research Agency in Applied Ecology, Wilshausen, France

ABSTRACT

Greylag goose populations are steadily increasing in north-western Europe. Although individuals breeding in the Netherlands have been considered mainly sedentary birds, those from Scandinavia or northern Germany fly towards their winter quarters, namely over France as far as Spain. This study aimed to determine the genetic structure of these birds, and to evaluate how goose populations mix. We used mitochondrial DNA and microsatellites from individuals distributed throughout the European Atlantic flyway, from breeding sites in Norway and the Netherlands to stopover and wintering sites in northern and south-western France. The mtDNA marker (CR1 D-Loop, 288 bp sequence, 144 ind.) showed 23 different haplotypes. The genetic distances amongst individuals sampled in Norway, northern France and the Netherlands were low (range 0.012–0.013). Individuals in south-western France showed a slightly higher genetic distance compared to all other sampling areas (ranges 0.018–0.022). The NJ tree does not show evidence of any single clades grouping together all individuals from the same geographic area. Besides, individuals from each site are found in different branches. Bayesian clustering procedures on 14 microsatellites (169 individuals) did not detect any geographically distinct cluster, and a high genetic admixture was recorded in all studied areas except for the individuals from the breeding sites in Norway, which were genetically very close. Estimation of migration rates through Bayesian inference confirms the scenario for the current mixing of goose populations.

Submitted 7 April 2015
Accepted 17 July 2015
Published 13 August 2015

Corresponding author
Marco Cucco, cucco@unipmn.it

Academic editor
Michael Wink

Additional Information and
Declarations can be found on
page 17

DOI [10.7717/peerj.1161](https://doi.org/10.7717/peerj.1161)

© Copyright
2015 Pellegrino et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Biodiversity, Zoology

Keywords mtDNA, Microsatellites, Greylag goose, Genetic structure, France and Norway

INTRODUCTION

The greylag goose (*Anser anser*) is widespread throughout the Palearctic. In Europe, the main breeding populations are located in central and northern countries, and the species rarely breeds in Mediterranean areas (*Cramp, 1977; Hagemeyer & Blair, 1997; BirdLife International, 2004*). European populations show different patterns of movement. Although individuals breeding in Scotland and the Netherlands are considered sedentary birds (*Delany & Scott, 2006*), those from Scandinavia or central Europe fly longer distances, namely over France to Spain, with some individuals reaching north Africa (*Fox et al., 2010; Nilsson et al., 2013*). Icelandic breeders winter in Ireland and Britain, and greylags from

Russia reach the regions bordering the eastern Mediterranean, Black and Caspian seas. Individuals with morphological characters ascribed to the oriental subspecies *rubrirostris* have been observed on rare occasions in western Europe (Cramp, 1977).

Widescale movement patterns have been studied through the recapture or resighting of marked birds (coloured neck collars and leg rings, see Nilsson *et al.*, 2013). These methods gave valuable information about the origin of birds that were found in moulting areas (Nilsson, Kahlert & Persson, 2001), flying, or staging in winter quarters. Birds from Sweden and Norway fly to Denmark and/or the Netherlands (SOVON, 1987; Persson, 1993; Andersson *et al.*, 2001). Recent monitoring of greylags tagged with GPS devices in Norway show that approximately 30–50% can stay in the Netherlands during the whole wintering season, whereas others migrate to France or Spain. These geese all return to their previous breeding sites, thus showing a high breeding site fidelity (Boos *et al.*, 2012; M Boos, 2014, unpublished data). According to Ramo *et al.* (2012), an increasing number of greylag geese winter at higher latitude. A noticeable effect of climatic changes probably explains this increasing tendency for geese to winter more closely to their breeding grounds.

The European Atlantic flyway covers a vast area stretching from northern France to Spain and Portugal, with arrivals from Scandinavia, Poland, Denmark and Germany (Fouquet, Schricke & Fouque, 2009). The situation in France is particularly complicated, because noticeable fluxes of geese coming from northern or Central Europe are found not only along the Atlantic flyway but also in other areas located in central and south-eastern France. The departure areas of these birds have yet to be fully determined, and the timing of migration can probably differ depending on the origin of the populations (Fouquet, 1991; Comolet-Tirman, 2009). Furthermore, the relative proportion of geese travelling to France and originating from different countries may change over time (Pistorius, Follestad & Taylor, 2006; Pistorius *et al.*, 2007). However, data from neck-collared or ringed geese can be skewed by variations in the marking and resighting efforts of the countries involved (Nilsson, 2007; Nilsson *et al.*, 2013), and this makes it difficult to fully define the composition of goose subpopulations migrating south from observational data alone.

Genetics have become a useful tool in the study of migration and wintering patterns. Recent studies on Anseriforms examined spatial structure along the flyways or in wintering zones, then compared it to genetic data in breeding areas. In the king eider (*Somateria spectabilis*), strong site fidelity to wintering areas and pair formation at wintering quarters indicated a population structure defined by wintering rather than nest-site philopatry (Pearce *et al.*, 2004). However, genetic analyses of mtDNA and microsatellite alleles showed a lack of spatial genetic structure, suggesting the possible existence of flows with homogenized gene frequencies. In the mallard (*Anas platyrhynchos*), single nucleotide markers were used to investigate population structure on a continental scale throughout the northern hemisphere. This genetic analysis found a general panmixia, suggesting that mallards form a single large, interbreeding population (Kraus *et al.*, 2013). The tufted duck (*Aythya fuligula*) shows high breeding site fidelity, but mtDNA and microsatellite markers revealed an extensive population admixture on the wintering ground (Liu, Keller & Heckel, 2012; Liu, Keller & Heckel, 2013). In the common pochard (*Aythya ferina*), genetic

differentiation was observed among Eurasian breeding populations, but no evidence of genetic structure was detected for pochards sampled on European wintering grounds (Liu, Keller & Heckel, 2011).

Relatively few studies have investigated the genetic aspects of European geese of the genus *Anser*, and the subject has not been thoroughly investigated at all in the greylag goose. Studies by Ruokonen *et al.* (2004), Ruokonen, Aarvak & Madsen (2005) examined the genetic variability in two species of conservation concern, the lesser white-fronted goose (*Anser erythropus*) and the pink-footed goose (*Anser brachyrhynchus*), and investigated the phylogenetic relationship between seven *Anser* species (Ruokonen, Kvist & Lumme, 2000). A small amount of genetic differentiation between species has been observed in this genus (Ruokonen, Kvist & Lumme, 2000; Johnsen *et al.*, 2010). Actually, mitochondrial DNA showed the presence of highly fragmented populations in two species of conservation concern, the lesser white-fronted goose (*Anser erythropus*, Ruokonen *et al.*, 2004) and the pink-footed goose (*Anser brachyrhynchus*, Ruokonen, Aarvak & Madsen, 2005).

However, population genetics among species of geese has not been investigated to date. Here we used both mitochondrial DNA and microsatellites to study the characteristics of greylag geese from two breeding areas (the Netherlands and north-western Norway) and two wintering zones (northern France and south-western France). This study investigates to what extent populations are genetically differentiated. Genetic structure could have been increased by the fragmentation of breeding geese in separated areas, or on the contrary, a limited genetic structure could have been developed by (i) the widespread practice of amateur breeding and selling of geese (Hagemeijer & Blair, 1997; Wang *et al.*, 2010), (ii) the recent increase in the size of several populations (Klok *et al.*, 2010), and (iii) the habit of European geese to rest several times during their flight toward their winter quarters (Fouquet, Schricke & Fouque, 2009) at stopover sites where individuals from distant areas can admix and form pair bonds.

Knowledge of the genetic structure and diversity of greylag goose populations is a necessary scientific basis to manage this emblematic species (Lorenz, 1966) and decide on appropriate action for its conservation (Kampe-Persson, 2002) in the light of serious recent conflicts with agricultural and habitat protection interests in most north European countries (Klok *et al.*, 2010).

METHODS

Sample collection and DNA extraction

We analyzed feather samples from 174 greylag geese (Appendix S1) collected over the European Atlantic flyway (from two breeding grounds: in north and western Norway, and six staging grounds: in the Netherlands, northern France and south-western France; see Table 1 and Fig. 1). One additional individual was collected in the Republic of Kalmykia in an area associated with the eastern *rubrirostris* subspecies (Cramp, 1977). During the 2010/2011 and 2011/2012 (including 1–10 February) wintering seasons in France, goose feathers were obtained from greylag geese collected during the legal hunting period in natural areas by hunters collaborating with the study. Samples from the Netherlands were

Table 1 Genetic variability of mtDNA CR1 D-Loop.

	Population	Season	N	Polymorphic sites	Hapl.	Private hapl.	<i>h</i>	π	<i>K</i>	<i>D</i>	<i>FS</i>
South-western France	Charente-Maritime	Autumn-winter	10	16	7	1	0.933 (0.06)	0.01790 (0.00254)	5.156	-0.409	-0.664
	Gironde	Autumn-winter	20	12	7	0	0.821 (0.06)	0.01519 (0.0012)	4.374	1.052	1.190
	Landes	Autumn-winter	36	19	13	4	0.798 (0.06)	0.0161 (0.0013)	4.633	0.0376	-0.360
North- France	Nord	Autumn-winter	17	10	7	0	0.833 (0.06)	0.01054 (0.0016)	3.025	0.014	0.433
	Oise	Autumn-winter	13	12	9	1	0.949 (0.042)	0.01309 (0.0020)	3.769	-0.103	-2.691*
Northern Europe	Norway Finnmark	Late spring	11	6	3	1	0.564 (0.13)	0.00947 (0.002)	2.727	1.31175	3.038
	Norway Vega	Late spring	23	7	3	0	0.170 (0.10)	0.0021 (0.0016)	0.609	-2.147*	0.270
	Netherlands	Autumn	14	11	8	2	0.890 (0.006)	0.01244 (0.0013)	3.571	0.541	-1.112
All Samples			144	20	23	-	0.823 (0.022)	0.01331 (0.0006)	3.819	-1.419	-4.515*

Notes.

h, haplotype diversity; π , nucleotide diversity; *D*, Tajima's *D*; *FS*, Fu's *FS*.

Standard errors are showed in brackets.

* $P < 0.05$.

obtained on wild free-ranging geese collected in natural areas in the Zeeland region (near Rilland) by a local hunter before the 20th of September in 2011 and 2013, i.e., before the arrival of geese breeding in Norway or in Sweden (Nilsson, 2007; M Boos, pers. obs., 2014 based on GPS data). Samples from Norway were obtained from birds that were collected during the spring and summer legal hunting seasons, or from geese that were caught during the moulting period in 2010 and 2011 by A.F. for the Nordic Greylag Goose Project, which studies the ecology of the Norwegian breeding goose population (Nilsson, 2007). Feather calami were stored in ethanol at -20°C , and total DNA was extracted using the commercial NucleoSpin[®] Tissue kit (Macherey–Nagel, Düren, Germany). After extraction, genomic DNA was stocked at -20°C .

Mitochondrial DNA sequencing

Partial mitochondrial control region (CR1 D-Loop 288 bp) was amplified in 144 of the 174 individuals (Appendix S1) using L180 (5'TGGTTATGCATATTCGTGCATAGA'3) and H466 (5'TTTCACGTGAGGAGTACGACTAAT'3) primers (Ruokonen, Kvist & Lumme, 2000). PCR amplifications were carried out in a Bio-Rad thermal cycler (Bio-Rad Laboratories Inc., Hercules, California, USA). PCR reaction was performed in a final volume of 25 μl containing 0.4 μl dNTPs (10 mM), 1 μl MgCl_2 (25 mM), 0.3 μl of each primer (25 pmol/ μl), 2.5 μl 10 \times buffer, 0.4 μl Taq polymerase (5 unit/ μl ; QIAGEN), ddH₂O and genomic DNA (20–100 ng/ μl). The selected cycling profile included a 4 min preliminary




Figure 1 Greylag goose distribution in Europe. Main *Anser anser* flyways from breeding (red) to wintering (blue) areas (modified from IUCN, 2015). Pie charts indicate the proportion of different haplotypes (mtDNA) found in each sampled population. Colours are identical to those used in the haplotype network (Fig. 2), and haplotypes found in one area alone are the same colour.

denaturation cycle at 94 °C followed by 32 denaturation, annealing and extension cycles (30 s at 94 °C, 30 s at 58 °C and 30 s at 72 °C, respectively) before a final extension of 7 min. Negative controls were included for amplification procedures to detect contaminations.

The PCR product was purified using the EXO-SAP procedure with Exonuclease I (Exo; Fermentas, Burlington, Canada) and Shrimp Alkaline Phosphatase (SAP; Fermentas, Burlington, Canada). The purification cycle consisted of 30 min at 37 °C, then 15 min at 80 °C to deactivate the enzymes followed by a 10 min cooling-down step at 4 °C. DNA concentration was determined after electrophoresis in 1.8% agarose gels (TBE 1%) stained with ethidium bromide and visualized in a UV-trans illuminator Gel Doc XR (Bio-Rad Laboratories Inc., Hercules, California, USA) using the Molecular Imager ChemiDoc XRS System and Quantity One software (Bio-Rad Laboratories Inc., Hercules, California, USA).

Sequencing was carried out at Macrogen Laboratories (Amsterdam, The Netherlands) in an ABI 3730xl Analyzer (Applied Biosystems).

Raw electropherograms were checked visually using FINCHTV (Geospiza Inc., Seattle, WA, USA; <http://www.geospiza.com>), and sequences were aligned with ClustalW algorithm in BIOEDIT 7.05 (Hall, 1999). The haplotype network was calculated in Network 4.6 (Fluxus Technology Ltd, Clare, Suffolk, England; fluxus-engineering.com) using the

median joining procedure (MJ: [Bandelt, Forster & Rohl, 1999](#)). DNASP version 5 ([Librado & Rozas, 2009](#)) was used to estimate mtDNA haplotype diversity (h), nucleotide diversity (π) and the mean number of pairwise differences (k) in the sampled areas. Demographic and/or spatial population expansion events were investigated using the mismatch distribution implemented in DNASP v. 5. MEGA 5.0 ([Tamura et al., 2011](#)) was used to perform the neighbour-joining method (NJ: [Saitou & Nei, 1987](#)), clustering pairwise Tamura-Nei's genetic distances between haplotypes (TN93: [Tamura & Nei, 1993](#)). Support for the internodes in the NJ tree was assessed by bootstrap percentages (BP: [Felsenstein, 1988](#)) after 1,000 resampling steps. One sequence of *Anser anser anser* (GenBank [AF159962](#)) from Finland and another of *Anser anser rubrirostris* (GenBank [AF159963](#)) from Slimbridge Wetland Center, England, were included as reference sequences in tree construction. A sequence of the lesser white-fronted goose (*Anser erythropus*, GenBank [AY072580](#)) and the bean goose (*Anser fabalis*, GenBank [AB551534](#)) were used as outgroups.

Maximum likelihood (ML) and maximum parsimony (MP) trees were obtained through the DNAML, CONSENSE, DNAPARS programmes in PHYLIP 3.67 ([Felsenstein, 2005](#)). Bootstrap values were based on 1,000 replicates, and the tree topologies were visualized with FIGTREE 1.3.1 ([Rambaut, 2008](#)). The best substitution model for molecular evolution was selected using the corrected Akaike Information Criterion (AICc, [Burnham & Anderson, 2004](#)) in jMODELTEST ([Posada, 2008](#)). Maximum likelihood bootstrap supports were estimated by performing 100 runs with 1,000 bootstrap replicates.

The partition of mtDNA diversity within and among the sampled geographical populations were investigated by running analyses of molecular variance (AMOVA, [Excoffier, Smouse & Quattro, 1992](#)) using ARLEQUIN 3.3 ([Excoffier & Lischer, 2010](#)).

Microsatellite genotyping

A total of 169 of the 174 samples ([Appendix S1](#)) were genotyped by PCR amplification at 14 microsatellite loci (Ans02, Ans04, Ans07, Ans13, Ans17, Ans18, Ans21, Ans24, Ans25, Aal μ 1b, Aph12b, Aph19b, Smo7b, Hhi μ 1b) that had previously been isolated and tested in *Anser anser* ([Weiß et al., 2008](#)). We used PCR reactions, thermal profiles, fluorescent dye and multiplex sets, as indicated by [Weiß et al. \(2008\)](#). Microsatellite genotyping was performed on an ABI Prism 3100 Genetic Analyzer (Applied Biosystems) using the Macrogen Inc. GenScan service (Seoul, Korea). Negative controls were included for amplification procedures. Results were analysed in GENEMAPPER v. 4.0 (Applied Biosystems, Foster City, California).

Allele frequencies, standard diversity indices, observed heterozygosity (H_O) and expected heterozygosity (H_E) for each locus and population were calculated in GENALEX v. 6 ([Peakall & Smouse, 2006](#)).

We performed a factorial correspondence analysis (FCA) of individual multilocus scores in GENETIX 4.05 ([Belkhir et al., 2004](#)) to describe genetic clusters.

GENEPOP 3.4 ([Raymond & Rousset, 1995; Rousset, 2008](#)) was used to calculate departures from the Hardy–Weinberg equilibrium (HWE) at each locus and within each population. Statistics were computed with Markov chain parameters at default settings.

We used ARLEQUIN 3.5 (Excoffier & Lischer, 2010) to estimate the genetic variance within and between populations through a hierarchical Analysis of Molecular Variance (AMOVA; Excoffier, Smouse & Quattro, 1992).

The genetic structure of the sampled populations was computed using Bayesian clustering procedures in STRUCTURE v. 2.3 (Pritchard, Stephens & Donnelly, 2000; Falush, Stephens & Pritchard, 2003), without prior information about the origin and under an admixed model. Analyses were performed where $K = 1-10$ with 50×10^5 iterations following a burn-in period of 50×10^4 iterations; all simulations were independently replicated four times for each K . We explored the optimal value of K by plotting the average estimated $\text{LnP}(D)$ (Ln probability of the data) and using ΔK statistics (Evanno, Regnaut & Goudet, 2005) calculated using STRUCTURE HARVESTER 0.6.93 (Earl & VonHoldt, 2012). CLUMPP v. 1.1.2 (Jakobsson & Rosenberg, 2007) and DISTRUCT v. 1.1 (Rosenberg, 2003) were used to align the cluster membership coefficients of the five STRUCTURE runs and display the results.

We investigated the presence of bottleneck events with BOTTLENECK v. 1.2.02 software (Cornuet & Luikart, 1996) for two models: the infinite alleles (IAM, Maruyama & Fuerst, 1985) and the two-phase model (TPM, Di Rienzo et al., 1994).

Migration rate was estimated using the Bayesian inference approach implemented in BAYESASS 3.0.3 (Wilson & Rannala, 2003). We performed 10 runs of 9×10^6 iterations with a burn-in of 10%, and a sampling frequency of 200. Delta values were varied for all parameters, and resulted in acceptance rates between 40% and 60% of the total iterations (Wilson & Rannala, 2003).

Finally, isolation by distance was tested via Mantel tests with GENEPOP (Raymond & Rousset, 1995; Rousset, 2008; see also Legendre & Fortin, 2010); F_{ST} and geographic distance were compared using 1,000 random permutations. The geographic distance connecting samples was represented by Euclidean (linear geographic) distances computed in QGIS (QGIS Development Team, 2014).

RESULTS

mtDNA

The mtDNA marker sequences (CR1 D-Loop 288 bp) showed 23 haplotypes defined by ten polymorphic sites and distributed in eight locations (Appendix S2). Among the 23 haplotypes found (GenBank accession numbers KT276333–KT276355), 14 haplotypes were shared by 2–47 individuals. According to the study areas, we found a total of nine private haplotypes, the majority of which occurred in the Landes (SW France) population (Table 1, Appendix S2). The diversity indices for mtDNA revealed moderate levels of genetic variation in the greylag goose in all sampled areas (Table 1). Haplotype diversity showed high values in all groups (range 0.798–0.949) except in breeding areas in Finnmark (0.564 ± 0.13 SD) and Vega, Norway (0.170 ± 0.10 SD).

The genetic distances recorded in Norway, northern France and the Netherlands were low (range 0.012–0.013). A slightly higher genetic distance was observed in south-west

Table 2 Tamura Nei genetic distance assessed by mtDNA. Below the diagonal the genetic distance values and above the diagonal their standard errors.

	SW France - Landes	SW France - Gironde	N France - Nord	N France Oise	Netherlands	Norway - Finnmark	Norway - Vega	SW France Charente M
SW France - Landes		0.004	0.004	0.004	0.004	0.005	0.006	0.005
SW France - Gironde	0.016		0.004	0.004	0.004	0.004	0.005	0.005
N France - Nord	0.015	0.014		0.004	0.004	0.004	0.004	0.005
N France Oise	0.016	0.015	0.013		0.004	0.004	0.004	0.005
Netherlands	0.015	0.014	0.013	0.013		0.004	0.004	0.005
Norway - Finnmark	0.018	0.016	0.014	0.013	0.014		0.003	0.006
Norway - Vega	0.017	0.014	0.012	0.012	0.012	0.008		0.006
SW France Charente M	0.020	0.019	0.019	0.018	0.018	0.022	0.019	

France in comparison to all other sampling sites (ranges 0.018–0.022), while the two breeding sites in Norway were genetically very close (Table 2).

The NJ tree shows that clades are composed of a wide variety of different geese from different areas. Individuals from each site were present in different branches (Fig. 2). Besides, none of the clades grouped together individuals originating from the same areas. About half the individuals were grouped together with the GenBank reference sequence relating to the *anser* subspecies, while the remainder were either grouped with the sequence relating to the *rubrirostris* subspecies or differed clearly from both subspecies. Very similar topologies were obtained from trees generated with other tree-building methods (MP and ML; not shown).

The haplotype median-joining network (Fig. 2) was concordant with the phylogenetic tree topology and did not reveal any geographic structures. The number of mutations separating the different haplotypes was low (max = 10).

Whilst 88.85% of the total genetic variance shown in hierarchical AMOVA was within populations, the remaining 11.15% occurred among populations. This indicates a small differentiation between the sampled areas.

Non-significant raggedness indices indicated a good fit to a model of population expansion in all sampled areas. Mismatch distribution results also suggested a population expansion in all areas except the Gironde region ($P = 0.044$) and Finnmark ($P = 0.042$) (Appendix S4). Fu's FS value (Table 1) was only significantly negative for the Oise region, and was consistent with a demographic expansion for all other areas.

Microsatellites

Among the 15 microsatellites previously isolated by Weiß *et al.* (2008) Ans26 was shown to be monomorphic in all investigated individuals. The remaining 14 polymorphic microsatellite loci showed 2–12 different alleles per locus ($n = 169$ individuals; Table 3).

Observed and expected heterozygosities were moderate, with similar values in each sampled population (H_o ranging from 0.374 to 0.484 and H_e from 0.433 to 0.549). Geese from the Landes wintering area exhibited the highest number of private alleles ($n = 3$, Appendix S3).




Figure 2 Network and NJ tree. (A) median-joining haplotype network. Areas of circles represent different sampled mtDNA haplotypes in proportion to their frequencies. Distances between haplotypes are proportional to the number of base differences. Colours match those utilized in Fig. 1, and haplotypes found in one area alone are the same colour. (B) neighbour-joining tree based on 280 bp of CR1. Sampled areas are labelled with abbreviations; NF, Northern France (Oise and Nord); SWF, South-Western France (Gironde, Charente Maritime and Landes); Nor, Norway; Neth, Netherlands. Numbers below branches indicate bootstrap values; only values above 50% are shown, most of the clades are supported by low bootstrap values.

Table 3 Summary of genetic variation at 14 microsatellite loci.

	Population	N	Na	Ne	Ho	He	HWE (P)
South-western France	Charente-Maritime	9	8.931 (0.071)	2.468 (0.340)	0.483 (0.082)	0.512 (0.051)	<0.001
	Gironde	24	4.286 (0.633)	2.575 (0.329)	0.462 (0.063)	0.504 (0.065)	<0.001
	Landes	45	4.643 (0.684)	2.625 (0.344)	0.484 (0.048)	0.542 (0.049)	<0.001
North France	Nord	17	4.000 (0.584)	2.680 (0.332)	0.483 (0.072)	0.549 (0.056)	<0.001
	Oise	15	3.857 (0.573)	2.702 (0.403)	0.385 (0.061)	0.527 (0.063)	<0.001
North Europe	Norway Finnmark	11	3.357 (0.372)	2.137 (0.266)	0.374 (0.072)	0.433 (0.065)	<0.01
	Norway Vega	34	4.571 (0.661)	2.485 (0.312)	0.476 (0.060)	0.529 (0.048)	<0.001
	Netherlands	14	3.929 (0.549)	2.352 (0.305)	0.447 (0.063)	0.493 (0.052)	<0.001
All samples		169	4.027 (0.200)	2.503 (0.115)	0.449 (0.023)	0.511 (0.020)	<0.001

Notes.

N, number of individuals; Na, No. of different alleles; Ne, No. of effective alleles; Ho, observed heterozygosity; He, expected heterozygosity.




Figure 3 Factorial correspondence analysis (FCA) of microsatellites data. Outliers concern individuals from Nord (4 ind.), Oise, Gironde and Finnmark (1 ind.) populations.

Genetic structure was visualized using factorial correspondence analysis (FCA) in each population (Fig. 3). The plot shows an absence of phylogeographic structure in the different investigated areas: individuals from different areas overlap, with the exclusion of four samples from Nord, one sample from Finnmark, one from Oise and one from Gironde.

Significant departures from HWE, due to heterozygote deficit and related to positive FIS values, were observed in all populations (Table 3, Appendix S3).

AMOVA analyses showed that 97.9% of the total genetic variance in geese was significantly distributed within populations ($p < 0.001$), while only 2.1% was distributed among populations. Overall fixation index F_{ST} from AMOVA was 0.02105, indicating a low differentiation between areas.

Table 4 Mean estimated number of migrants between breeding and wintering sites as calculated with BayesAss (standard deviations in parentheses). Values on the diagonal (in bold) represent the estimated proportion of resident individuals in each population.

Migration from	Migration into							
	Netherlands	NF-Nord	NF-Oise	NO-Finnmark	NO-Vega	SWF-Charente Maritime	SWF-Gironde	SWF-Landes
Netherlands	0.8904 (0.0338)	0.0167 (0.0160)	0.0151 (0.0147)	0.0152 (0.0144)	0.0160 (0.0154)	0.0152 (0.0146)	0.0157 (0.0151)	0.0157 (0.0148)
NF- Nord	0.0145 (0.0139)	0.9009 (0.0316)	0.0135 (0.0130)	0.0140 (0.0135)	0.0144 (0.0138)	0.0133 (0.0129)	0.0149 (0.0143)	0.0145 (0.0141)
NF- Oise	0.0142 (0.0135)	0.0460 (0.0238)	0.6820 (0.0146)	0.0144 (0.0139)	0.0138 (0.0132)	0.0136 (0.0132)	0.2023 (0.0344)	0.0137 (0.0133)
NO- Finnmark	0.0191 (0.0183)	0.0212 (0.0198)	0.0174 (0.0165)	0.8619 (0.0394)	0.0215 (0.0210)	0.0177 (0.0164)	0.0219 (0.0211)	0.0194 (0.0184)
NO-Vega	0.0085 (0.0081)	0.0078 (0.0078)	0.0078 (0.0078)	0.0081 (0.0079)	0.9439 (0.0192)	0.0079 (0.0077)	0.0081 (0.0079)	0.0081 (0.0077)
SWF- Charente Maritime	0.2072 (0.0379)	0.0175 (0.0167)	0.0173 (0.0165)	0.0176 (0.0166)	0.0181 (0.0174)	0.6872 (0.0192)	0.0177 (0.0168)	0.0174 (0.0166)
SWF- Gironde	0.0114 (0.0112)	0.0107 (0.0105)	0.0111 (0.0102)	0.0199 (0.0160)	0.0109 (0.0104)	0.0110 (0.0107)	0.9142 (0.0270)	0.0109 (0.0105)
SWF- Landes	0.0069 (0.0069)	0.0068 (0.0067)	0.0063 (0.0063)	0.0067 (0.0066)	0.0065 (0.0064)	0.0063 (0.0063)	0.0068 (0.0067)	0.9538 (0.0164)

STRUCTURE analyses, performed without the use of prior information on sample locations, showed a maximum ΔK at $K = 4$, while likelihood values reached a plateau at $K = 7$ (Fig. 4). Graphs show no evidence of phylogeographic structure across sampled populations, whatever the K value. With $K = 4$, only 23 individuals with individual q_i values were each assigned to a single cluster: two individuals from Finnmark, three from Vega and one from Gironde were attributed to cluster 1; one individual from Netherland, two from Finnmark, six from Vega, one from Charente Maritime, two from Gironde and one from Landes were attributed to cluster 2; two individuals from Oise and two from Landes were assigned to cluster 3. All other birds had a highly mixed genotype. In the case of $K = 7$, five other individuals, one from Finnmark, Charente Maritime and Gironde and two from Vega, were assigned to the same cluster with $q_i > 0.90$.

Bottleneck events tested under IAM revealed a significant excess of heterozygotes (evidence of a recent bottleneck) in Nord, Landes and Oise populations (Wilcoxon sign-rank tests, all $P < 0.05$). Analysis under TPM only confirmed a recent bottleneck event for the Nord population ($P < 0.05$).

BAYESAss detected a low migration rate among localities and a high proportion of local individuals ($>68\%$, Table 4), suggesting that the flows among different areas were limited. Indeed, the analysis found a high proportion of local geese in six populations ($>90\%$). In two cases, gene flow appears to be strongly asymmetrical, with many birds moving from Charente Maritime to the Netherlands ($20.7\% \pm 3.79$ SD) and from Oise to Gironde ($20.2\% \pm 3.44$ SD), but not in the opposite direction (1.5% and 1.1% respectively).




Figure 4 Structure analysis. Estimated population structure in Greylag Goose sampled populations. Each vertical line represents one individual and each colour represents a single cluster.

The Mantel test calculated on geographic and genetic distances yielded a non-significant correlation coefficient ($r = 0.107$; $P = 0.08$), suggesting that there is no strong relationship between geographic and genetic distances.

DISCUSSION

In this study we used a pool of 14 microsatellites isolated by [Weiß et al. \(2008\)](#) for greylag goose parentage in the long-established goose population at Konrad Lorenz Research Station, Grünau, Austria ([Lorenz, 1966](#); [Hirschenhauser, Möstl & Kotrschal, 1999](#)). We found that these microsatellites can be successfully employed for geese sampled in a wide range of localities along the European Atlantic flyway. This is the first large scale

study showing a moderate genetic variability of mtDNA and nuclear DNA in all French wintering areas and in the Netherlands, with slightly lower mtDNA variability in the Norwegian breeding sites. A moderate genetic variability in the greylag goose was already reported two decades ago by *Blaakmeer (1995)*, and has been found in other species of geese (*Anser erythropus*: *Ruokonen et al., 2004*; *Ruokonen et al., 2010*; *Anser brachyrhynchus*: *Ruokonen, Aarvak & Madsen, 2005*). Low genetic variability also seems to be typical for other Anatidae species (*Aythya ferina*: *Liu, Keller & Heckel, 2011*; *Aythya fuligula*: *Liu, Keller & Heckel, 2012*). Interestingly, our results show that the genotypes deviated from Hardy–Weinberg expectation at eight loci, and in all study areas deviation was due to heterozygote deficiency. Additionally, the deficit of heterozygotes matched with positive FIS values. These results could be related to different factors such as population substructuring or recent population growth (*Cornuet & Luikart, 1996*).

Genetic distances between the different areas were low (range from 0.012 to 0.017) and the hierarchical AMOVA showed genetic variance to mainly occur within populations. These findings could be explained by a small differentiation between the sampled areas and a general admixture of greylag goose populations in our western European study region. However, it should be taken into account that genetic divergences in geese are characteristically very small, with the lowest interspecific divergence reported here for avian species (*Ruokonen, Kvist & Lumme, 2000*). The genetic tree shows that different branches include individuals from each sampling area. No single branch exclusively grouped together individuals originating from the same zone. Moreover, birds sampled in the western part of the breeding range, traditionally ascribed to the *anser* subspecies, were not separated from birds collected in the eastern part that were traditionally assigned to the *rubrirostris* subspecies (*Kampe-Persson, 2002*). Birds from Iceland, Scotland and coastal Norway have been sometimes separated as a race, *sylvestris*, classified in the *anser* group (*Snow, Perrins & Cramp, 1998*). Although Icelandic and Scottish birds were absent from this study, individuals from the Norwegian west coast did not appear to be clearly distinct from other European geese. Our present results slightly differ from the findings of *Blaakmeer's (1995)* preliminary study, which reported genetic differences between breeders in two Dutch sites in comparison to breeding sites in south Sweden and Norway. However, *Blaakmeer's* analyses show significant differences for only one of six minisatellites, in only two of the three Dutch areas studied.

Interestingly, the ANS19 sequence was recently found in the white Roman goose in Taiwan. This race is widely bred for commercial purposes, and has been found to originate from the European species (*Anser anser*, *Wang et al., 2010*). Our data confirm the presence of this sequence in Europe, particularly in the breeding population of the Norwegian west coast.

Haplotypes ANS02, ANS08, ANS11, ANS14 and ANS23 were only found during the winter in France, and were absent in Norway and the Netherlands: this result could indicate that some of the geese arriving in France came from areas we did not sample on the breeding grounds. Ring recoveries and resighting records indicate that these birds probably originated from northern Germany, Poland, Denmark and Sweden (*Nilsson et al., 2013*).

The haplotype network confirmed the tree configuration. There was no geographic pattern, and the number of mutations separating the different nodes was very low. This confirms the low genetic distance between our studied populations in the large north-western European population (as defined by [Delany & Scott, 2006](#)), and may reflect the rapid population expansion ([Aris-Brosou & Excoffier, 1996](#)).

Data obtained from nuclear DNA by microsatellites were in accordance with findings from mtDNA. As the mtDNA is uniparentally inherited whereas microsatellites are part of the biparentally inherited nuclear DNA, a difference between the two genomes would have indicated the presence of sex-biased dispersal ([Fahey, Ricklefs & Dewoody, 2014](#)). However, sex-biased dispersal seems to be unlikely in greylag geese for three reasons: the family unit remains together at least until autumn migration, the birds tend to pair before returning to the breeding grounds, and males and females have long-term pair bonds ([Rohwer & Anderson, 1988](#); [Doherty et al., 2002](#)). Sex-biased dispersal in birds is probably not a species constant ([Clarke, Saether & Roskaft, 1997](#)). Within Anatidae in general, sex-biased dispersal was not detected in several species ([Doherty et al., 2002](#); [Mabry et al., 2013](#)), while it was found in some species such as the white-fronted goose ([Anser erythropus, Ruokonen et al., 2010](#)), the common eider ([Somateria mollissima, Paulus & Tiedemann, 2003](#)), and the spectacled eider ([Somateria fisheri, Scribner et al., 2001](#)).

In our study of microsatellites, individuals from different geographic localities were found to be combined in the Factorial Correspondence Analysis representation. Bayesian structure analysis resulted in a best combination of four or seven groups, according to ΔK and $\text{LnP}(D)$ methods respectively. As seen in our previous analyses, no geographic clustering was observed inside these STRUCTURE groups. Almost all individuals, with few exceptions, showed admixed genotype regardless of the number of groups considered in the analysis.

The high mixing of genotypes and the lack of geographic structure among our studied populations could be interpreted in the light of the data obtained through ringing activity and the extensive neck-banding programme carried out in Scandinavia from 1984 to 2004 ([Nilsson, 2007](#); [Voslamber, Knecht & Kleijn, 2010](#)). Ring recoveries and visual observations showed that Scandinavian geese breeding in different zones can admix not only in the moulting areas ([Nilsson, Kahlert & Persson, 2001](#)) but also along the European Atlantic flyway, i.e., in the Netherlands ([Voslamber, Knecht & Kleijn, 2010](#)), France ([Fouquet, Schricke & Fouquet, 2009](#); [Nilsson et al., 2013](#)) and Spain ([Ramo et al., 2012](#)) where they can form pairbonds. Besides this Scandinavian data, the monitoring of collared and/or ringed individuals performed in other European areas showed the presence of birds in France originating from Germany, the Czech Republic and Poland. Populations that breed further east do not seem to reach France in winter ([Kampe-Persson, 2010](#)). From these data it appears that the greylag geese that cross France or winter there could result from a mixture of populations from different areas.

Our findings are somewhat unexpected if one assumes that the fragmentation of breeding populations into separate areas during the first part of the last century ([Hagemeyer & Blair, 1997](#); [Kampe-Persson, 2002](#)), should have led to an increase in genetic

structure. Moreover, birds breeding in the Netherlands have recently become highly sedentary (Fox *et al.*, 2010), and this may also have contributed to the increase in genetic structure (Blaakmeer, 1995). However, a genetic panmixia could have been promoted by the widespread amateur breeding and selling of geese, and the recent increase and dispersal of several wild goose populations (Klok *et al.*, 2010). In particular, geese with pink bills and legs, most probably *rubrirostris* subspecies, have been spreading in Europe over the last few decades; their natural flyway toward wintering areas crosses other European countries (from Russia to Hungary, the Balkan States and Italy) but does not reach France.

The breeding of geese is a widespread practice among amateurs, who can easily obtain both goslings and adults with a grey wild appearance (B and G Vaschetti, pers. comm., 2014). In some cases geese were released as part of assisted restoring projects and are now indistinguishable from the wild individuals (Kampe-Persson, 2010). Besides, birds with white plumage are common in breeding farms. In Asia, white geese are mostly descendants of the swan goose (*Anser cygnoides*). Even if descendants of *Anser anser* can also be found there, they are usually farmed in Europe (Wang *et al.*, 2010). Although the two species can hybridize in captivity, hybrids can be detected through karyotype (Shahin, Ata & Shnaf, 2014) or genotype examinations (Sun *et al.*, 2014). The contribution of escaped white form geese to the admixture observed in wild populations is probably low given the high assortative mating of wild greylag geese, their long-term monogamous pair bonds, female-bonded clan structure, long parent-offspring relationships, and elaborate patterns of mutual social support (Hirschenhauser *et al.*, 2000; Kotrschal, Scheiber & Hirschenhauser, 2010).

Our findings on the greylag goose genetic admixture are similar to those reported in the snow goose (*Chen caerulescens*, Avise *et al.*, 1992) and the barnacle goose (*Branta leucopsis*, Jonker *et al.*, 2013). Despite the high rate of site philopatry seen in the snow goose, which has also shown a high increase in population over the last decades, mtDNA markers showed no clear distinctions between nesting populations across species range (Avise *et al.*, 1992). The barnacle goose recently changed its migratory traditions, and new populations differing in migratory distance were observed. Genetic data showed an admixture between all populations, despite the assumed traditions of migration within areas and the presence of a newly established nonmigratory population in the Netherlands (Jonker *et al.*, 2013). A lack of genetic structure in wintering areas was also found in four species of Anatidae, namely the common pochard, the mallard, the king eider and the tufted duck (Pearce *et al.*, 2004; Liu, Keller & Heckel, 2011; Liu, Keller & Heckel, 2012; Kraus *et al.*, 2013; Liu *et al.*, 2013), and in the black-tailed godwit (*Limosa limosa*, Lopes *et al.*, 2013). The mixing of breeding populations in wintering areas is believed to be a common phenomenon in birds, because the breeding ranges of most species are considerably larger than their wintering ranges (Winker & Graves, 2008). However, migratory populations vary in the degree to which individuals from distinct breeding localities mix on different sites. Therefore, to understand population demographics and genetic diversification, it is crucial to pinpoint which populations mix on breeding and wintering grounds (Chabot *et al.*, 2012).

Our DNA-based estimates of migration during the wintering period indicated a low rate of exchange between our sampled areas. In five of eight areas the vast majority of

individuals (86–95%) did not switch among the different zones, and a moderate exchange (about 20%) was only observed from Charente Maritime to the Netherlands and from Oise to Gironde. These results seem to support the hypothesis that the French wintering birds arrive from various areas, including zones that are not sampled here (i.e., Germany, Poland), while the contribution of the Norwegian population represents only a portion of the whole assemblage (Fouquet, Schricke & Fouque, 2009). This low exchange rate is also supported by evidence of great changes in spatial ecology recorded in 28 GPS tagged western European greylag geese, i.e., very small home ranges on wintering areas ($<8.9 \pm 2.5 \text{ km}^2$) compared to 2–5 fold values obtained in staging areas during migratory and premigratory periods (M Boos, 2014, unpublished data).

Our data show evidence of genetic bottlenecks in just three groups under IAM, all located along the same flyway (Nord, Oise and Landes), and in a single case (Nord) under TPM. The discrepancy between IAM and TPM could be related to the different ability of the mutation models to detect bottleneck events. Empirical data suggest that TPM is the most appropriate model for microsatellite loci (Ellegren, 2000; Ellegren, 2004) while IAM results should be interpreted with caution (Cornuet & Luikart, 1996). We did not observe any sign of bottlenecks in the breeding populations: this indicates that greylag geese have never suffered any severe demographic reduction, even at the beginning of the past century when the number of breeding individuals was low in several European areas (Kampe-Persson, 2002).

Greylag goose populations are steadily increasing in north-western Europe (Kampe-Persson, 2002). The large number of geese in some areas is now in conflict with agricultural interests, since geese not only forage in natural environments but also forage on crop fields, and claims for a need to control the species are widespread (Klok et al., 2010). Our data suggest that the migratory geese harvested over France (about 20,000 geese/year, see Landry & Migot, 2000) show a relatively high diversity of origin. From this result it is difficult to conclude if there is a strong impact on a specific breeding population. Future studies could analyze other European breeding and wintering areas; this could clarify the status of the different populations and subspecies on the continent (the main *Anser anser anser* and *A. a. rubrirostris*, as well as the *sylvestris* forms from Iceland, Scotland and Norway), and help to build an effective international management strategy for this migratory species (Chabot et al., 2012).

ACKNOWLEDGEMENTS

Our thanks to all the hunters who provided feather samples from France, to Wilmer Remijnse who provided feather samples from Netherlands, and to Paul Shimmings for his field assistance in Norway. We are grateful to Bruno and Gabriella Vaschetti who provided useful information on goose farming, the International Union for Conservation of Nature (IUCN) which provided an updated distribution map, and to Joanna Lignot-Munro for language editing.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The study was supported by the French National Fund for Biological Research in Wildlife Species to Mathieu Boos. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
French National Fund.

Competing Interests

Arne Follestad is an employee of the Department of Terrestrial Ecology, Norsk Institutt for Naturforskning and Mathieu Boos is an employee of Naturaconst@, Research Agency in Applied Ecology.

Author Contributions

- Irene Pellegrino conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Marco Cucco conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Arne Follestad reviewed drafts of the paper, field work, collecting samples.
- Mathieu Boos conceived and designed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:
GenBank accession numbers: [KT276333](#) to [KT276355](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1161#supplemental-information>.

REFERENCES

- Andersson Å, Follestad A, Nilsson L, Persson H. 2001. Migration patterns of Nordic greylag geese *Anser anser*. *Ornis Svecica* 11:19–58.
- Aris-Brosou S, Excoffier L. 1996. The impact of population expansion and mutation rate heterogeneity on DNA sequence polymorphism. *Molecular Biology and Evolution* 13:494–504 DOI [10.1093/oxfordjournals.molbev.a025610](https://doi.org/10.1093/oxfordjournals.molbev.a025610).
- Avise JC, Alisauskas RT, Nelson WS, Ankney CD. 1992. Matriarchal population genetic structure in an avian species with female natal philopatry. *Evolution* 46:1084–1096 DOI [10.2307/2409758](https://doi.org/10.2307/2409758).

- Bandelt H-J, Forster P, Rohlf A. 1999.** Median-Joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16**:37–48 DOI [10.1093/oxfordjournals.molbev.a026036](https://doi.org/10.1093/oxfordjournals.molbev.a026036).
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F. 2004.** GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Montpellier: Laboratoire Genome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II.
- BirdLife International. 2004.** *Birds in Europe, population estimates, trends and conservation status*. Birdlife International edition. Cambridge.
- Blaakmeer K. 1995.** *Genetic variation within the west European population of the Greylag goose (Anser anser)*. Groningen: University of Groningen.
- Boos M, Schricke V, Green AJ, Shimmings P, Lefranc H, Amat JA, Ramo C, Follestad A. 2012.** Migration of greylag geese tagged in Norway and Spain using GPS devices: first results from a new joint European Research Program. In: *14th meeting of the goose specialist group*, vol. 8. Steinkjer, Norway.
- Burnham KP, Anderson DR. 2004.** Multimodel inference: understanding AIC and BIC in model selection. *Sociological Methods & Research* **33**:261–304 DOI [10.1177/0049124104268644](https://doi.org/10.1177/0049124104268644).
- Chabot AA, Hobson KA, Van Wilgenburg SL, McQuat GJ, Lougheed SC. 2012.** Advances in linking wintering migrant birds to their breeding-ground origins using combined analyses of genetic and stable isotope markers. *PLoS ONE* **7**:e43627 DOI [10.1371/journal.pone.0043627](https://doi.org/10.1371/journal.pone.0043627).
- Clarke AL, Saether B-E, Roskaft E. 1997.** Sex biases in avian dispersal: a reappraisal. *Oikos* **79**:429–438 DOI [10.2307/3546885](https://doi.org/10.2307/3546885).
- Comolet-Tirman J. 2009.** *L'Oie cendrée Anser anser en France et en Europe. Dynamique de population, statuts de conservation, voies de migration et dates de migration pré-nuptiale*. Service du Patrimoine Naturel Département Ecologie et Gestion de la Biodiversité.
- Cornuet JM, Luikart G. 1996.** Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* **144**:2001–2014.
- Cramp S. 1977.** *Handbook of the birds of Europe, the Middle East, and North Africa, The birds of the Western Palearctic*, Vol. 1. Oxford: Oxford University Press.
- Delany S, Scott D. 2006.** *Waterbird population estimates*. Fourth Edition. Wageningen: Wetland International.
- Di Rienzo A, Peterson AC, Garza JC, Valdes AM, Slatkin M, Freimer NB. 1994.** Mutational processes of simple-sequence repeat loci in human populations. *Proceedings of the National Academy of Sciences of the United States of America* **91**:3166–3170 DOI [10.1073/pnas.91.8.3166](https://doi.org/10.1073/pnas.91.8.3166).
- Doherty PF, Nichols JD, Tautin J, Voelzer JF, Smith GW, Benning DS, Bentley VR, Bidwell JK, Bollinger KS, Brazda AR, Buelna EK, Goldsberry JR, King RJ, Roetker FH, Solberg JW, Thorpe PP, Wortham JS, Service USG, Wildlife P, Road BF. 2002.** Sources of variation in breeding-ground fidelity of mallards (*Anas platyrhynchos*). *Behavioral Ecology* **13**:543–550 DOI [10.1093/beheco/13.4.543](https://doi.org/10.1093/beheco/13.4.543).
- Earl DA, VonHoldt BM. 2012.** STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* **4**:359–361 DOI [10.1007/s12686-011-9548-7](https://doi.org/10.1007/s12686-011-9548-7).
- Ellegren H. 2000.** Microsatellite mutations in the germline: implications for evolutionary inference. *Trends in Genetics* **16**:551–558 DOI [10.1016/S0168-9525\(00\)02139-9](https://doi.org/10.1016/S0168-9525(00)02139-9).
- Ellegren H. 2004.** Microsatellites: simple sequences with complex evolution. *Nature Reviews. Genetics* **5**:435–445 DOI [10.1038/nrg1348](https://doi.org/10.1038/nrg1348).
- Evanno G, Regnaut S, Goudet J. 2005.** Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* **14**:2611–2620 DOI [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x).

- Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**:564–567 DOI [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x).
- Excoffier L, Smouse PE, Quattro JM. 1992.** Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**:479–491.
- Fahey AL, Ricklefs RE, Dewoody JA. 2014.** DNA-based approaches for evaluating historical demography in terrestrial vertebrates. *Biological Journal of the Linnean Society* **112**:367–386 DOI [10.1111/bij.12259](https://doi.org/10.1111/bij.12259).
- Falush D, Stephens M, Pritchard JK. 2003.** Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**:1567–1587.
- Felsenstein J. 1988.** Phylogenies from molecular sequences: inference and reliability. *Annual Review of Genetics* **22**:521–565 DOI [10.1146/annurev.ge.22.120188.002513](https://doi.org/10.1146/annurev.ge.22.120188.002513).
- Felsenstein J. 2005.** *PHYLIP (Phylogeny Inference Package)*. Version 3.6.
- Fouquet M. 1991.** Migration et hivernage de l'oie cendrée (*Anser anser*) en France. Rôle et importance du Centre-Ouest. *Oiseau et la Revue Française d'Ornithologie* **61**:111–130.
- Fouquet M, Schricke V, Fouque C. 2009.** Greylag geese *Anser anser* depart earlier in spring: an analysis of goose migration from western France over the years 1980–2005. *Wildfowl* **59**:143–151.
- Fox AD, Ebbinge BS, Mitchell C, Heinicke T, Aarvak T, Colhoun K, Clausen P, Dereliev S, Faragó S, Koffijberg K, Kruckenberg H, Pihl S, Jeugd H. 2010.** Current estimates of goose population sizes in western Europe, a gap analysis and an assessment of trends. *Ornis Svecica* **20**:115–127.
- Hagemeijer EJM, Blair MJ. 1997.** *The EBCC Atlas of European breeding birds: their distribution and abundance*. London: T. & A. D. Poyser.
- Hall T. 1999.** BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**:95–98.
- Hirschenhauser K, Mo E, Bernard A, John W, Kurt D. 2000.** Endocrine and behavioural responses of male greylag geese *Anser anser* to pairbond challenges during the reproductive season. *Ethology* **106**:63–77 DOI [10.1046/j.1439-0310.2000.00499.x](https://doi.org/10.1046/j.1439-0310.2000.00499.x).
- Hirschenhauser K, Möstl E, Kotschal K. 1999.** Seasonal patterns of sex steroids determined from feces in different social categories of greylag geese (*Anser anser*). *General and Comparative Endocrinology* **114**:67–79 DOI [10.1006/gcen.1998.7236](https://doi.org/10.1006/gcen.1998.7236).
- IUCN. 2015.** *The IUCN red list of threatened species*. Version 2015.1. Available at <http://www.iucnredlist.org> (accessed on 11 May 2015).
- Jakobsson M, Rosenberg NA. 2007.** CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**:1801–1806 DOI [10.1093/bioinformatics/btm233](https://doi.org/10.1093/bioinformatics/btm233).
- Johnsen A, Rindal E, Ericson PGP, Zuccon D, Kerr KCR, Stoeckle MY, Lifjeld JT. 2010.** DNA barcoding of Scandinavian birds reveals divergent lineages in trans-Atlantic species. *Journal of Ornithology* **151**:565–578 DOI [10.1007/s10336-009-0490-3](https://doi.org/10.1007/s10336-009-0490-3).
- Jonker RM, Kraus RHS, Zhang Q, Van Hooft P, Larsson K, Van der Jeugd HP, Kurvers RHJM, Van Wieren SE, Loonen MJJE, Crooijmans RPMA, Ydenberg RC, Groenen MAM, Prins HHT. 2013.** Genetic consequences of breaking migratory traditions in barnacle geese *Branta leucopsis*. *Molecular Ecology* **22**:5835–5847 DOI [10.1111/mec.12548](https://doi.org/10.1111/mec.12548).

- Kampe-Persson H. 2002.** *Anser anser* greylag goose. *BWP Update* **4**:181–216.
- Kampe-Persson H. 2010.** Naturalised geese in Europe. *Ornis Svecica* **20**:155–173.
- Klok C, Schekkerman H, Willems F, Ebbinge B, Van Turnhout C. 2010.** Analysis of population development and effectiveness of management in resident greylag geese *Anser anser* in the Netherlands. *Animal Biology* **60**:373–393 DOI [10.1163/157075610X523260](https://doi.org/10.1163/157075610X523260).
- Kotrschal K, Scheiber IB, Hirschenhauser K. 2010.** Individual performance in complex social systems: the greylag goose example. In: Kappeler P, ed. *Animal behaviour: evolution and mechanisms*. Berlin: Springer Verlag, 121–148.
- Kraus RHS, Van Hooft P, Megens H-J, Tsvey A, Fokin SY, Ydenberg RC, Prins HHT. 2013.** Global lack of flyway structure in a cosmopolitan bird revealed by a genome wide survey of single nucleotide polymorphisms. *Molecular Ecology* **22**:41–55 DOI [10.1111/mec.12098](https://doi.org/10.1111/mec.12098).
- Landry P, Migot P. 2000.** Enquête nationale sur les tableaux de chasse à tir: saison 1998/1999. *Faune Sauvage Cahiers Techniques* **251**:8–17.
- Legendre P, Fortin M-J. 2010.** Comparison of the Mantel test and alternative approaches for detecting complex multivariate relationships in the spatial analysis of genetic data. *Molecular Ecology Resources* **10**:831–844 DOI [10.1111/j.1755-0998.2010.02866.x](https://doi.org/10.1111/j.1755-0998.2010.02866.x).
- Librado P, Rozas J. 2009.** DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **25**:1451–1452 DOI [10.1093/bioinformatics/btp187](https://doi.org/10.1093/bioinformatics/btp187).
- Liu Y, Keller I, Heckel G. 2011.** Range-wide genetic population structure of common pochard (*Aythya ferina*): a potentially important vector of highly pathogenic avian influenza viruses. *Ecology and Evolution* **1**:529–545 DOI [10.1002/ece3.46](https://doi.org/10.1002/ece3.46).
- Liu Y, Keller I, Heckel G. 2012.** Breeding site fidelity and winter admixture in a long-distance migrant, the tufted duck (*Aythya fuligula*). *Heredity* **109**:108–116 DOI [10.1038/hdy.2012.19](https://doi.org/10.1038/hdy.2012.19).
- Liu Y, Keller I, Heckel G. 2013.** Temporal genetic structure and relatedness in the Tufted Duck *Aythya fuligula* suggests limited kin association in winter. *Ibis* **155**:499–507 DOI [10.1111/ibi.12059](https://doi.org/10.1111/ibi.12059).
- Liu S, Li Y, Lu J, Su X, Tang M, Zhang R, Zhou L, Zhou C, Yang Q, Ji Y, Yu DW, Zhou X. 2013.** SOAPBarcode: revealing arthropod biodiversity through assembly of Illumina shotgun sequences of PCR amplicons. *Methods in Ecology and Evolution* **4**:1142–1150 DOI [10.1111/2041-210X.12120](https://doi.org/10.1111/2041-210X.12120).
- Lopes RJ, Alves JA, Gill JA, Gunnarsson TG, Hooijmeijer JCEW, Lourenco PM, Masero JA, Piersma T, Potts PM, Rabacal B, Reis S, Sanchez-Guzman JM, Santiago-Quesada F, Villegas A. 2013.** Do different subspecies of Black-tailed Godwit *Limosa limosa* overlap in Iberian wintering and staging areas? Validation with genetic markers. *Journal of Ornithology* **154**:35–40 DOI [10.1007/s10336-012-0865-8](https://doi.org/10.1007/s10336-012-0865-8).
- Lorenz KZ. 1966.** The triumph ceremony of the greylag goose, *Anser anser*. *Philosophical Transactions of the Royal Society B: Biological Sciences* **251**:477 DOI [10.1098/rstb.1966.0038](https://doi.org/10.1098/rstb.1966.0038).
- Mabry KE, Shelley EL, Davis KE, Blumstein DT, Van Vuren DH. 2013.** Social mating system and sex-biased dispersal in mammals and birds: a phylogenetic analysis. *PLoS ONE* **8**:e57980 DOI [10.1371/journal.pone.0057980](https://doi.org/10.1371/journal.pone.0057980).
- Maruyama T, Fuerst PA. 1985.** Population bottlenecks and non equilibrium models in population genetics. II. Number of alleles in a small population that was formed by a recent bottleneck. *Genetics* **111**:675–689.
- Nilsson L. 2007.** The Nordic greylag goose (*Anser anser*) project. *Aves* **44**:177–184.

- Nilsson L, Follestad A, Guillemain M, Schricke V, Voslamber B. 2013. France as a staging and wintering area for greylag geese *Anser anser*. *Wildfowl* 63:24–39.
- Nilsson L, Kahlert J, Persson H. 2001. Moults and moult migration of greylag geese *Anser anser* from a population in Scania, south Sweden. *Bird Study* 48:129–138
DOI 10.1080/00063650109461211.
- Paulus KB, Tiedemann R. 2003. Ten polymorphic autosomal microsatellite loci for the Eider duck *Somateria mollissima* and their cross-species applicability among waterfowl species (Anatidae). *Molecular Ecology Notes* 3:250–252 DOI 10.1046/j.1471-8286.2003.00414.x.
- Peakall R, Smouse PE. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295
DOI 10.1111/j.1471-8286.2005.01155.x.
- Pearce JM, Talbot SL, Pierson BJ, Petersen MR, Scribner KT, Dickson DL, Mosbech A. 2004. Lack of spatial genetic structure among nesting and wintering King Eiders. *The Condor* 106:229–240 DOI 10.1650/7357.
- Persson H. 1993. Arrival patterns of greylag geese *Anser anser* in the Guadalquivir Marismas. *Wildfowl* 44:19–23.
- Pistorius PA, Follestad A, Nilsson L, Taylor FE. 2007. A demographic comparison of two Nordic populations of greylag geese *Anser anser*. *Ibis* 149:553–563
DOI 10.1111/j.1474-919X.2007.00682.x.
- Pistorius PA, Follestad A, Taylor FE. 2006. Declining winter survival and fitness implications associated with latitudinal distribution in Norwegian greylag geese *Anser anser*. *Ibis* 148:114–125 DOI 10.1111/j.1474-919X.2006.00498.x.
- Posada D. 2008. jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* 25:1253–1256 DOI 10.1093/molbev/msn083.
- Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- QGIS Development Team. 2014. QGIS geographic information system. In: *Open Source Geospatial Foundation Project*. Available at <http://qgis.osgeo.org>.
- Rambaut A. 2008. *FigTree: Tree Figure drawing tool 2006–2009*. Version 1.3.1. Institute of Evolutionary Biology, University of Edinburgh. Available at <http://tree.bio.ed.ac.uk>.
- Ramo C, Amat JA, Calderón J, Gómez-Crespo E, Navedo JG, Green AJ, Jubete F, Masero JA, Palacios J, Rodríguez-Alonso M, Boos M, Schricke V. 2012. Distribution and population trends of wintering greylag geese in Spain. In: *14th Meeting of the Goose Specialist Group*, vol. 28. Steinkjer: Nord-Trøndelag University College.
- Raymond M, Rousset F. 1995. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86:248–249.
- Rohwer FC, Anderson MG. 1988. Female-biased philopatry, monogamy, and the timing of pair formation in migratory waterfowl. In: Johnston E, Richard F, eds. *Current Ornithology v.5*. Boston: Springer, 187–221.
- Rosenberg NA. 2003. Distruct: a program for the graphical display of population structure. *Molecular Ecology Notes* 4:137–138 DOI 10.1046/j.1471-8286.2003.00566.x.
- Rousset F. 2008. GENEPOP'007: a complete reimplementations of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8:103–106
DOI 10.1111/j.1471-8286.2007.01931.x.

- Ruokonen M, Aarvak T, Chesser RK, Lundqvist A-C, Merilä J. 2010.** Temporal increase in mtDNA diversity in a declining population. *Molecular Ecology* **19**:2408–2417.
- Ruokonen M, Aarvak T, Madsen J. 2005.** Colonization history of the high-arctic pink-footed goose *Anser brachyrhynchus*. *Molecular Ecology* **14**:171–178
DOI [10.1111/j.1365-294X.2004.02380.x](https://doi.org/10.1111/j.1365-294X.2004.02380.x).
- Ruokonen M, Kvist L, Aarvak T, Markkola J, Morozov V, Øien IJ, Syroechkovsky Jr EE, Tolvanen P, Lumme J. 2004.** Population genetic structure and conservation of the lesser white-fronted goose *Anser erythropus*. *Conservation Genetics* **5**:501–512
DOI [10.1023/B:COGE.0000041019.27119.b4](https://doi.org/10.1023/B:COGE.0000041019.27119.b4).
- Ruokonen M, Kvist L, Lumme J. 2000.** Close relatedness between mitochondrial DNA from seven *Anser* goose species. *Journal of Evolutionary Biology* **13**:532–540
DOI [10.1046/j.1420-9101.2000.00184.x](https://doi.org/10.1046/j.1420-9101.2000.00184.x).
- Saitou N, Nei M. 1987.** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**:406–425.
- Scribner KT, Petersen MR, Fields RL, Talbot SL, Pearce JM, Chesser RK. 2001.** Sex-biased gene flow in Spectacled Eiders (Anatidae): inferences from molecular markers with contrasting modes of inheritance. *Evolution* **55**:2105–2115 DOI [10.1111/j.0014-3820.2001.tb01325.x](https://doi.org/10.1111/j.0014-3820.2001.tb01325.x).
- Shahin AAB, Ata ATM, Shnaf ASMA. 2014.** Karyotype and C-banding pattern of the domestic geese *Anser anser* populations (Aves: Anatidae) in Egypt. *Folia Biologica* **62**:49–58.
- Snow DW, Perrins CM, Cramp S. 1998.** *The complete birds of the western Palaearctic: On CD-ROM*. Oxford: Oxford University Press.
- SOVON. 1987.** *Atlas van de Nederlandse vogels*. Nederland, Leiden: SOVON Vogelonderzoek Nederland, Arnhem NL.
- Sun J, Zhang S, He D-Q, Chen S-Y, Duan Z-Y, Yao Y-G, Liu Y-P. 2014.** Matrilineal genetic structure of domestic geese. *Journal of Poultry Science* **51**:130–137.
- Tamura K, Nei M. 1993.** Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**:512–526.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011.** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**:2731–2739
DOI [10.1093/molbev/msr121](https://doi.org/10.1093/molbev/msr121).
- Voslamber B, Knecht E, Kleijn D. 2010.** Dutch greylag geese *Anser anser*: migrants or residents? *Ornis Svecica* **20**:207–214.
- Wang CM, Way TD, Chang YC, Yen NT, Hu CL, Nien PC, Jea YS, Chen LR, Kao JY. 2010.** The origin of the white Roman goose. *Biochemical Genetics* **48**:938–943
DOI [10.1007/s10528-010-9374-8](https://doi.org/10.1007/s10528-010-9374-8).
- Weiβ BM, Poggemann K, Olek K, Foerster K, Hirschenhauser K. 2008.** Isolation and characterization of microsatellite marker loci in the greylag goose (*Anser anser*). *Molecular Ecology Resources* **8**:1411–1413 DOI [10.1111/j.1755-0998.2008.02339.x](https://doi.org/10.1111/j.1755-0998.2008.02339.x).
- Wilson GA, Rannala B. 2003.** Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* **163**:1177–1191.
- Winker K, Graves GR. 2008.** Genetic structure of breeding and wintering populations of Swainson's Warbler. *Wilson Journal of Ornithology* **120**:433–445 DOI [10.1676/07-073.1](https://doi.org/10.1676/07-073.1).

Appendix 1 - List of samples used in this study, with ID, population, sample provenance, collecting date, haplotype assignment and whether the sample was genotyped with microsatellites (STRs).

Sample	ID	POP	pop_name	Country	Locality	Sampling date	MtDNA Haplotype	STRs
aa68	PB/87	NL	Netherlands	Netherlands	Netherlands	13/10/11	ANS15	
aa69	PB/80	NL	Netherlands	Netherlands	Netherlands	13/10/11	ANS5	x
aa70	PB/86	NL	Netherlands	Netherlands	Netherlands	30/09/11	ANS3	x
aa71	PB/75	NL	Netherlands	Netherlands	Netherlands	15/09/11	ANS15	x
aa72	PB/81	NL	Netherlands	Netherlands	Netherlands	30/09/11	ANS3	x
aa73	PB/79	NL	Netherlands	Netherlands	Netherlands	30/09/11	ANS6	x
aa74	PB/85	NL	Netherlands	Netherlands	Netherlands	13/10/11	ANS3	x
aa75	PB/76	NL	Netherlands	Netherlands	Netherlands	27/09/11	ANS1	x
aa76	PB/78	NL	Netherlands	Netherlands	Netherlands	27/09/11	ANS7	x
aa77	PB/82	NL	Netherlands	Netherlands	Netherlands	27/09/11	ANS1	x
aa78	PB/88	NL	Netherlands	Netherlands	Netherlands	27/09/11	ANS3	x
aa79	PB/77	NL	Netherlands	Netherlands	Netherlands	30/09/11	ANS10	x
aa80	PB/83	NL	Netherlands	Netherlands	Netherlands	20/09/11	ANS1	x
aa81	PB/74	NL	Netherlands	Netherlands	Netherlands	15/09/11		x
aa82	PB/84	NL	Netherlands	Netherlands	Netherlands	20/09/11	ANS9	x
aa26	59/33	59/62	Nord	Northern France	Nord	2011/2012	ANS1	x
aa37	59/40	59/62	Nord	Northern France	Nord	2011/2012	ANS13	x
aa41	59/44	59/62	Nord	Northern France	Nord	2011/2012		x
aa43	59/32	59/62	Nord	Northern France	Nord	2011/2012	ANS1	x
aa48	59/31	59/62	Nord	Northern France	Nord			x
aa53	59/45	59/62	Nord	Northern France	Nord		ANS1	
aa58	59/34	59/62	Nord	Northern France	Nord		ANS1	x
aa67	59/30	59/62	Nord	Northern France	Nord	end Jan 2012		x
aa103	59/89	59/62	Nord	Northern France	Nord	15/10/11	ANS4	x
aa104	59/90	59/62	Nord	Northern France	Nord		ANS15	
aa105	59/91	59/62	Nord	Northern France	Nord	15/10/11	ANS15	
aa106	59/92	59/62	Nord	Northern France	Nord		ANS1	x
aa107	59/93	59/62	Nord	Northern France	Nord	12/11/11	ANS3	x
aa108	59/94	59/62	Nord	Northern France	Nord	12/11/11	ANS3	x
aa109	59/95	59/62	Nord	Northern France	Nord	12/11/11	ANS10	x
aa110	59/96	59/62	Nord	Northern France	Nord	12/11/11	ANS3	x
aa111	59/97	59/62	Nord	Northern France	Nord	06/11/11	ANS3	x

aa112	59/98	59/62	Nord	Northern France	Nord	24/10/11	ANS10	x
aa113	59/99	59/62	Nord	Northern France	Nord	24/10/11	ANS10	x
aa49	60/43	60/76/80	Oise	Northern France	Oise		ANS3	x
aa93	60/60	60/76/80	Oise	Northern France	Oise	10/02/12		x
aa94	60/55	60/76/80	Oise	Northern France	Oise	10/02/12	ANS10	x
aa95	60/54	60/76/80	Oise	Northern France	Oise	10/02/12	ANS4	x
aa96	60/59	60/76/80	Oise	Northern France	Oise	04/02/12	ANS12	x
aa97	60/58	60/76/80	Oise	Northern France	Oise	10/02/12		x
aa98	60/56	60/76/80	Oise	Northern France	Oise	10/02/12	ANS4	x
aa114	60/151	60/76/80	Oise	Northern France	Oise		ANS7	x
aa115	60/152	60/76/80	Oise	Northern France	Oise		ANS1	x
aa116	60/153	60/76/80	Oise	Northern France	Oise		ANS18	x
aa117	60/154	60/76/80	Oise	Northern France	Oise	28/11/10	ANS13	x
aa118	60/155	60/76/80	Oise	Northern France	Oise	28/11/10	ANS13	x
aa56	62/35	59/62	Nord	Northern France	Pas-de-Calais			x
aa62	62/36	59/62	Nord	Northern France	Pas-de-Calais		ANS7	x
aa119	FRA124	60/76/80	Oise	Northern France	Seine-Maritime		ANS3	x
aa45	80/37	60/76/80	Oise	Northern France	Somme	2011/2012	ANS6	
aa51	80/38	60/76/80	Oise	Northern France	Somme		ANS10	x
aa120	Fin100	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa121	Fin101	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa122	Fin102	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa123	Fin103	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa124	Fin104	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS18	X
aa126	Fin106	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa127	Fin107	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa128	Fin108	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS3	x
aa129	Fin109	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS19	x
aa130	Fin110	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS18	x
aa131	Fin111	Finm	Finnmark	Norway	Finnmark	01/06/10	ANS18	x
aa132	Veg113	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa133	Veg116	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa134	Veg123	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa135	Veg128	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa136	Veg131	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa137	Veg132	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa138	Veg134	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa139	Veg135	Norw	Vega	Norway	Vega	01/06/10		x

aa140	Veg137	Norw	Vega	Norway	Vega	01/06/10		x
aa141	Veg140	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa142	Veg141	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa143	Smo112	Norw	Vega	Norway	Vega	01/06/10		x
aa144	Smo114	Norw	Vega	Norway	Vega	01/06/10		x
aa145	Smo115	Norw	Vega	Norway	Vega	01/06/10		x
aa146	Smo117	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa147	Smo118	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa148	Smo119	Norw	Vega	Norway	Vega	01/06/10		x
aa149	Smo120	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa150	Smo121	Norw	Vega	Norway	Vega	01/06/10		x
aa151	Smo122	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa152	Smo125	Norw	Vega	Norway	Vega	01/06/10		x
aa153	Smo126	Norw	Vega	Norway	Vega	01/06/10		x
aa154	Smo127	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa155	Smo129	Norw	Vega	Norway	Vega	01/06/10		x
aa156	Smo130	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa157	Smo133	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa158	Smo136	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa159	Smo138	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa160	Smo139	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa161	Smo142	Norw	Vega	Norway	Vega	01/06/10		x
aa162	Smo146	Norw	Vega	Norway	Vega	01/06/10	ANS15	x
aa163	Smo147	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa164	No143	Norw	Vega	Norway	Vega	01/06/10	ANS3	x
aa165	No149	Norw	Vega	Norway	Vega	01/06/10	ANS16	x
aa166	17/160	17	Charente Maritime	South-western France	Charente Maritime	10/02/12	ANS3	x
aa167	17/161	17	Charente Maritime	South-western France	Charente Maritime	07/02/12	ANS7	x
aa168	17/162	17	Charente Maritime	South-western France	Charente Maritime	10/02/12	ANS15	x
aa169	17/163	17	Charente Maritime	South-western France	Charente Maritime	07/02/12	ANS15	x
aa170	17/164	17	Charente Maritime	South-western France	Charente Maritime	07/02/12	ANS20	
aa171	17/165	17	Charente Maritime	South-western France	Charente Maritime	07/02/12	ANS21	x
aa172	17/166	17	Charente Maritime	South-western France	Charente Maritime	07/02/12	ANS21	x
aa173	17/167	17	Charente Maritime	South-western France	Charente Maritime	10/02/12	ANS1	x
aa174	17/168	17	Charente Maritime	South-western France	Charente Maritime	10/02/12	ANS22	x
aa175	17/169	17	Charente Maritime	South-western France	Charente Maritime	05/02/12	ANS3	x
aa16	33/09	33	Gironde	South-western France	Gironde	end Jan 2012	ANS4	x
aa17	33/05	33	Gironde	South-western France	Gironde	end Jan 2012		x

aa18	33/11	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa19	33/12	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa22	33/25	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa23	33/03	33	Gironde	South-western France	Gironde	end Jan 2012		x
aa25	33/18	33	Gironde	South-western France	Gironde	08/11/11	ANS15	
aa27	33/08	33	Gironde	South-western France	Gironde	end Jan 2012	ANS3	x
aa28	33/02	33	Gironde	South-western France	Gironde	end Jan 2012	ANS7	x
aa29	33/07	33	Gironde	South-western France	Gironde	end Jan 2012	ANS16	x
aa30	33/17	33	Gironde	South-western France	Gironde	end Jan 2012		x
aa31	33/04	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa32	33/19	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa33	33/06	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa36	33/14	33	Gironde	South-western France	Gironde	end Jan 2012	ANS3	x
aa38	33/13	33	Gironde	South-western France	Gironde	end Jan 2012	ANS7	x
aa39	33/20	33	Gironde	South-western France	Gironde	end Jan 2012	ANS17	x
aa40	33/26	33	Gironde	South-western France	Gironde	end Jan 2012	ANS15	x
aa42	33/27	33	Gironde	South-western France	Gironde	end Jan 2012	ANS1	x
aa44	33/21	33	Gironde	South-western France	Gironde	end Jan 2012	ANS3	x
aa59	33/29	33	Gironde	South-western France	Gironde	end Jan 2012		x
aa60	33/23	33	Gironde	South-western France	Gironde	end Jan 2012	ANS15	x
aa64	33/24	33	Gironde	South-western France	Gironde	end Jan 2012	ANS3	x
aa65	33/16	33	Gironde	South-western France	Gironde	end Jan 2012		x
aa66	33/22	33	Gironde	South-western France	Gironde	end Jan 2012	ANS15	x
aa24	40/47	40/64	Landes	South-western France	Landes	08/11/11	ANS15	x
aa34	X/53	40/64	Landes	South-western France	Landes	2011/2012	ANS17	x
aa35	X/50	40/64	Landes	South-western France	Landes	2011/2012	ANS1	x
aa46	40/46	40/64	Landes	South-western France	Landes	14/12/11	ANS3	x
aa52	X/39	40/64	Landes	South-western France	Landes			x
aa54	X/49	40/64	Landes	South-western France	Landes	–	ANS13	x
aa55	40/42	40/64	Landes	South-western France	Landes	–	ANS15	x
aa61	X/41	40/64	Landes	South-western France	Landes	–	ANS15	x
aa83	40/73	40/64	Landes	South-western France	Landes	16/11/11	ANS1	x
aa84	40/69	40/64	Landes	South-western France	Landes	16/11/11	ANS3	x
aa85	40/61	40/64	Landes	South-western France	Landes	29/01/12	ANS11	x
aa86	40/65	40/64	Landes	South-western France	Landes	16/11/11		x
aa87	40/71	40/64	Landes	South-western France	Landes	16/11/11		x
aa89	40/70	40/64	Landes	South-western France	Landes	16/11/11	ANS1	x
aa90	40/72	40/64	Landes	South-western France	Landes	16/11/11	ANS15	

aa91	40/67	40/64	Landes	South-western France	Landes	04/02/12		x
aa100	X/66	40/64	Landes	South-western France	Landes	-		x
aa101	X/62	40/64	Landes	South-western France	Landes	-		x
aa102	X/63	40/64	Landes	South-western France	Landes	-		x
aa176	40/170	40/64	Landes	South-western France	Landes	02/02/12	ANS21	x
aa178	40/172	40/64	Landes	South-western France	Landes	09/02/12	ANS1	x
aa180	40/174	40/64	Landes	South-western France	Landes	04/02/12	ANS23	x
aa181	40/175	40/64	Landes	South-western France	Landes	09/02/12	ANS15	x
aa182	40/176	40/64	Landes	South-western France	Landes	02/02/12	ANS1	x
aa183	40/177	40/64	Landes	South-western France	Landes	01/02/12	ANS3	x
aa184	40/178	40/64	Landes	South-western France	Landes	01/02/12	ANS1	x
aa185	40/179	40/64	Landes	South-western France	Landes	03/02/12	ANS1	x
aa186	40/180	40/64	Landes	South-western France	Landes	03/02/12	ANS14	x
aa187	40/181	40/64	Landes	South-western France	Landes	10/02/12	ANS20	x
aa188	40/182	40/64	Landes	South-western France	Landes	10/02/12	ANS7	x
aa189	40/183	40/64	Landes	South-western France	Landes	12/02/12	ANS14	x
aa190	40/184	40/64	Landes	South-western France	Landes	12/02/12	ANS1	x
aa40-1		40/64	Landes	South-western France	Landes	-	ANS1	x
aa40-2		40/64	Landes	South-western France	Landes	-	ANS1	x
aa40-3		40/64	Landes	South-western France	Landes	-	ANS1	x
aa40-4		40/64	Landes	South-western France	Landes	-	ANS2	x
aa40-5		40/64	Landes	South-western France	Landes	-	ANS8	x
aa40-6		40/64	Landes	South-western France	Landes	-	ANS1	X
aa40-7		40/64	Landes	South-western France	Landes	-	ANS1	X
aa40-8		40/64	Landes	South-western France	Landes	-	ANS1	X
aa47	64/52	40/64	Landes	South-western France	Pyrenee Atlantique	-	ANS15	x
aa50	64/48	40/64	Landes	South-western France	Pyrenee Atlantique	-	ANS1	x
aa57	64/51	40/64	Landes	South-western France	Pyrenee Atlantique	-	ANS3	x
rub1				Russia	Kalmykia		17	

	France					North Europe			Total
	Charente Maritime	Gironde	Landes	Nord	Oise	Norway Finnmark	Norway Vega	Netherlands	
ANS1	1	7	15	5	1			3	32
ANS2			1						1
ANS3	2	4	4	4	2	7	21	4	48
ANS4		1		1	2				4
ANS5								1	1
ANS6					1			1	2
ANS7	1	2	1	1	1			1	7
ANS8			1						1
ANS9								1	1
ANS10				3	2			1	6
ANS11			1						1
ANS12					1				1
ANS13			1	1	2				4
ANS14			2						2
ANS15	2	4	6	2			1	2	17
ANS16		1					1		2
ANS17		1	1						2
ANS18					1	3			4
ANS19						1			1
ANS20	1		1						2
ANS21	2		1						3
ANS22	1								1
ANS23			1						1
Total	10	20	36	17	13	11	23	14	144

Appendix 3 – Summary of genetic variation at 14 microsatellite loci in sampled populations. Na = No. of different alleles; Ne = No. of effective alleles. Ho = observed heterozygosity; He = expected heterozygosity; F = fixation index; HWE = Hardy-Weinberg equilibrium.


Pop	Locus	N	Na	Ne	Ho	He	F
Netherlands	Ans02	13	9.000	4.390	0.692	0.772	0.103
	Ans04	14	3.000	2.947	0.214	0.661	0.676
	Ans07	14	3.000	1.782	0.286	0.439	0.349
	Ans17	14	7.000	4.667	0.643	0.786	0.182
	Ans18	14	4.000	2.190	0.214	0.543	0.606
	Ans21	14	3.000	1.640	0.500	0.390	-0.281
	Ans24	14	3.000	1.338	0.286	0.253	-0.131
	Ans25	14	6.000	3.769	0.786	0.735	-0.069
	Aalu1b	14	4.000	2.190	0.429	0.543	0.211
	Aph12	14	2.000	1.600	0.214	0.375	0.429
	Aph19	14	2.000	1.774	0.500	0.436	-0.146
	Smo7	14	2.000	1.508	0.286	0.337	0.152
	Hhiu1b	14	3.000	1.156	0.929	0.135	-5.868
	Ans13	14	4.000	1.980	0.286	0.495	0.423
Nord	Ans02	16	8.000	3.606	0.750	0.723	-0.038
	Ans04	17	4.000	2.934	0.706	0.659	-0.071
	Ans07	17	3.000	2.181	0.294	0.542	0.457
	Ans17	17	8.000	5.026	0.706	0.801	0.119
	Ans18	16	3.000	1.724	0.125	0.420	0.702
	Ans21	17	3.000	1.847	0.235	0.458	0.487
	Ans24	17	1.000	1.000	0.000	0.000	-
	Ans25	17	7.000	4.983	0.882	0.799	-0.104
	Aalu1b	17	4.000	3.778	0.824	0.735	-0.120
	Aph12	15	3.000	2.103	0.533	0.524	-0.017
	Aph19	17	2.000	1.973	0.412	0.493	0.165
	Smo7	17	2.000	1.710	0.412	0.415	0.008
	Hhiu1b	17	4.000	1.889	0.529	0.471	-0.125
	Ans13	17	4.000	2.766	0.353	0.638	0.447
Oise	Ans02	14	9.000	6.759	0.857	0.852	-0.006
	Ans04	15	4.000	3.147	0.267	0.682	0.609
	Ans07	14	2.000	1.912	0.214	0.477	0.551
	Ans17	15	6.000	4.206	0.467	0.762	0.388
	Ans18	14	6.000	2.667	0.357	0.625	0.429
	Ans21	15	3.000	1.779	0.200	0.438	0.543
	Ans24	15	2.000	1.069	0.067	0.064	-0.034
	Ans25	15	6.000	3.947	0.600	0.747	0.196
	Aalu1b	14	4.000	2.947	0.643	0.661	0.027
	Aph12	14	3.000	2.074	0.286	0.518	0.448
	Aph19	15	2.000	1.867	0.333	0.464	0.282
	Smo7	15	2.000	1.642	0.533	0.391	-0.364
	Hhiu1b	15	2.000	1.069	0.067	0.064	-0.034
	Ans13	14	3.000	2.741	0.500	0.635	0.213

Finnmark	Ans02	9	6.000	3.857	0.778	0.741	-0.050
	Ans04	10	4.000	2.597	0.400	0.615	0.350
	Ans07	10	3.000	2.151	0.200	0.535	0.626
	Ans17	11	5.000	3.270	0.818	0.694	-0.179
	Ans18	8	2.000	1.280	0.250	0.219	-0.143
	Ans21	11	3.000	2.283	0.364	0.562	0.353
	Ans24	10	1.000	1.000	0.000	0.000	-
	Ans25	10	4.000	2.667	0.700	0.625	-0.120
	Aalu1b	11	4.000	1.330	0.182	0.248	0.267
	Aph12	11	3.000	1.582	0.455	0.368	-0.236
	Aph19	11	3.000	1.603	0.273	0.376	0.275
	Smo7	11	2.000	1.198	0.000	0.165	1.000
	Hhiu1b	11	2.000	1.198	0.182	0.165	-0.100
	Ans13	11	5.000	3.903	0.636	0.744	0.144
Vega	Ans02	31	11.000	3.979	0.774	0.749	-0.034
	Ans04	34	4.000	3.129	0.382	0.680	0.438
	Ans07	31	5.000	2.023	0.355	0.506	0.298
	Ans17	34	5.000	2.919	0.588	0.657	0.105
	Ans18	33	6.000	2.209	0.273	0.547	0.502
	Ans21	31	3.000	1.795	0.613	0.443	-0.384
	Ans24	30	3.000	1.224	0.067	0.183	0.635
	Ans25	34	8.000	5.639	0.882	0.823	-0.073
	Aalu1b	34	4.000	2.305	0.588	0.566	-0.039
	Aph12	34	3.000	2.070	0.412	0.517	0.203
	Aph19	34	2.000	1.895	0.588	0.472	-0.245
	Smo7	34	2.000	1.524	0.206	0.344	0.401
	Hhiu1b	34	3.000	1.433	0.353	0.302	-0.167
	Ans13	34	5.000	2.645	0.588	0.622	0.054
Charente-Maritime	Ans02	9	8.000	5.400	0.889	0.815	-0.091
	Ans04	9	4.000	2.746	0.111	0.636	0.825
	Ans07	9	3.000	1.588	0.000	0.370	1.000
	Ans17	9	5.000	3.057	0.444	0.673	0.339
	Ans18	9	3.000	2.793	0.222	0.642	0.654
	Ans21	9	3.000	1.976	0.667	0.494	-0.350
	Ans24	9	2.000	1.385	0.333	0.278	-0.200
	Ans25	8	6.000	4.923	0.875	0.797	-0.098
	Aalu1b	9	4.000	2.282	0.556	0.562	0.011
	Aph12	9	2.000	1.800	0.778	0.444	-0.750
	Aph19	9	2.000	1.800	0.222	0.444	0.500
	Smo7	9	2.000	1.385	0.889	0.278	-2.200
	Hhiu1b	9	3.000	1.256	0.222	0.204	-0.091
	Ans13	9	3.000	2.160	0.556	0.537	-0.034
Gironde	Ans02	23	9.000	5.186	0.826	0.807	-0.023
	Ans04	24	4.000	3.892	0.333	0.743	0.551
	Ans07	24	4.000	1.466	0.208	0.318	0.344
	Ans17	24	7.000	3.959	0.542	0.747	0.275
	Ans18	23	5.000	1.853	0.217	0.460	0.528
	Ans21	23	3.000	1.772	0.348	0.436	0.202
	Ans24	23	1.000	1.000	0.000	0.000	-


	Ans25	24	8.000	4.645	0.750	0.785	0.044	
	Aalu1b	24	5.000	2.743	0.667	0.635	-0.049	
	Aph12	24	2.000	1.946	0.667	0.486	-0.371	
	Aph19	23	2.000	1.830	0.435	0.454	0.042	
	Smo7	24	2.000	1.180	0.583	0.153	-2.818	
	Hhiu1b	24	4.000	1.550	0.333	0.355	0.061	
	Ans13	23	4.000	3.032	0.565	0.670	0.157	
Landes	Ans02	45	11.000	4.500	0.756	0.778	0.029	
	Ans04	44	4.000	3.741	0.477	0.733	0.349	
	Ans07	44	3.000	1.955	0.227	0.488	0.535	
	Ans17	45	7.000	4.787	0.689	0.791	0.129	
	Ans18	45	6.000	1.777	0.333	0.437	0.238	
	Ans21	45	3.000	1.874	0.467	0.466	-0.001	
	Ans24	45	3.000	1.280	0.222	0.219	-0.015	
	Ans25	45	8.000	4.350	0.800	0.770	-0.039	
	Aalu1b	45	5.000	3.061	0.489	0.673	0.274	
	Aph12	45	2.000	1.776	0.378	0.437	0.135	
	Aph19	44	3.000	1.890	0.477	0.471	-0.014	
	Smo7	45	2.000	1.557	0.556	0.358	-0.553	
	Hhiu1b	45	4.000	1.529	0.356	0.346	-0.028	
	Ans13	45	4.000	2.675	0.556	0.626	0.113	

		N	Na	Ne	Ho	He	F	HWE	
								P-value	(se)
Mean for each locus (all populations confounded)	Ans02	20.000	8.875	4.710	0.790	0.780	-0,014	0.3783	(0.0209)
	Ans04	20.875	3.875	3.142	0.361	0.676	0,466	0.0000	(0.0000)
	Ans07	20.375	3.250	1.882	0.223	0.459	0,520	0.0000	(0.0000)
	Ans17	21.125	6.250	3.986	0.612	0.739	0,170	0.0000	(0.0000)
	Ans18	20.250	4.375	2.062	0.249	0.487	0,439	0.0000	(0.0000)
	Ans21	20.625	3.000	1.871	0.424	0.461	0,071	0.1042	(0.003)
	Ans24	20.375	2.000	1.162	0.122	0.125	0,051	0.0558	(0.0022)
	Ans25	20.875	6.625	4.365	0.784	0.760	-0,033	0.3506	(0.0153)
	Aalu1b	21.000	4.250	2.579	0.547	0.578	0,073	0.0218	(0.0022)
	Aph12	20.750	2.500	1.869	0.465	0.459	-0,020	0.0406	(0.0015)
	Aph19	20.875	2.250	1.829	0.405	0.451	0,107	0.1316	(0.0038)
	Smo7	21.125	2.000	1.463	0.433	0.305	-0,547	0.0003	(0.0001)
	Hhiu1b	21.125	3.125	1.385	0.371	0.255	-0,794	0.0707	(0.0034)
	Ans13	20.875	4.000	2.738	0.505	0.621	0,190	0.0004	(0.0002)


Appendix 4 - Mismatch distribution calculated for all individuals and each population.




All Samples




Charente Maritime




Gironde




Landes




Nord




Oise



Norway Vega



Norway Finnmark



Netherlands